

Dermil

INJECTABLE

Antiinflammatory, Antiallergic.
Sterile aqueous suspension
for canines and felines



FORMULA: Triamcinolone acetonide: 0.6 g. Excipients c.s. 100 ml. / **INSTRUCTIONS:** Skin diseases such as allergy, dermatitis, pruritus, eczema, urticaria. Respiratory diseases of allergic origin. For all types of osteoarticular inflammatory process, arthritis, bursitis, tenosynovitis, myositis, rheumatoid syndrome. **SUPPLY AND DOSAGE:** Intramuscular, subcutaneous, intra-articular, intrasynovial. Canines: intramuscular or subcutaneous via: 0.1 to 0.2 mg./Kg. (0.33 ml. per each 10 kg.). Felines: intramuscular or subcutaneous via: 0.1 to 0.2 mg./kg. (0.15 ml. per animal). Infiltrations: Canines and felines 1.2 to 1.8 mg. (0.2 to 0.3 ml.) / **PRESENTATION:** Case containing 3 ampoules

- For the treatment of allergies, urticaria and dermatitis
- One single application for entire treatment

Dermil

CREAM

Analgesic, Antiinflammatory, Antibiotic,
Antifungal, Anesthetic, Antihistamine
for canines and felines



FORMULA: Betamethasone 17 valerate: 0.10 g.; Gentamycin sulfate: 0.10 g.; Clotrimazole: 1 g.; Vitamin A: 0.6g.; Lidocaine: 0.6g.; Chlorhexidine digluconate (20%): 5g.; Excipients c.s. 100 g. **INSTRUCTIONS:** Allergy, urticaria, dermatitis of allergic origin, localized, wet or dry eczema, conditions of allergic origin and other diseases of the skin that require treatment with corticosteroids. As an antiinflammatory and restorative of the skin. **SUPPLY AND DOSAGE:** External use, apply locally enough on the affected area, 2 to 3 times a day. Cover to avoid licking. / **PRESENTATION:** Pot of 50 gr.

- The most complete FORMULA in market
- Easy absorption
- With the addition of Vitamin A

Dermil

TABLETS

Antihistamine-Antiinflammatory
for canines and felines



FORMULA: Triamcinolone: 1.25 mg.; Chlorpheniramine maleate: 2.1 mg.; Vitamin "A": 5000 U.I., Excipients c.s. 1 tablet. / **INSTRUCTIONS:** Allergy, urticaria, dermatitis of allergic origin, pruritus (widespread or localized), eczema (wet or dry). As an antiinflammatory and restorative of the skin. / **SUPPLY AND DOSAGE:** Orally. 1 tablet per each 20 kg. liveweight. / **PRESENTATION:** Package of 100 and 400 tablets.

- For the treatment of allergies, urticaria and dermatitis.
- 1 tablet per each 20 kg.
- With Vitamin "A"
- Unique in market

Novul

TABLETS

Hormonal, contraceptive
for canines and felines



FORMULA: Chlormadinone acetate: 2 mg.; excipients c.s. 1 tablet. / **INSTRUCTIONS:** Interruption of estrus. Postpone estrus, metrorrhagia, nymphomania, alterations of the ovarian cycle, pseudopregnancy and hyper-sexuality in the male. Prevention of estrus in female canines and felines, in anestrus. / **SUPPLY AND DOSAGE:** Orally. To interrupt the estrus: 1 tablet per each 5 kg. liveweight for 8-10 days. To postpone estrus: 1 tablet per each 10 kg. liveweight for 4 weeks. Nymphomania, pseudopregnancy and hypersexuality in the male: 1 tablet per each 5 kg. liveweight according to effects. / **PRESENTATION:** Pack of 100 and 400

- For interruption or postponement of estrus

Novul

INJECTABLE

Hormonal, contraceptive
for canines and felines



FORMULA: Medroxyprogesterone acetate 50 mg.; Excipients c.s. 1 ml / **INSTRUCTIONS:** Prevention of estrus in female canines and felines, in anestrus. **SUPPLY AND DOSAGE:** Subcutaneous via. Canines and felines: 1 ml. (50 mg.) per animal, during anestrus. Repeat every 5/6 months. **PRESENTATION:** Case containing 12 ampoules of 1 ml. each one.

- Injectable contraceptive of choice.
- Only one application prevents estrus.

Doloprofen 100mg.

TABLETS

CARPROPHENE
Non-steroidal
anti-inflammatory drug
for oral use for dogs



FORMULA: Carprofeno 100 mg., Excipients csp 1 tablet. / **INSTRUCTIONS:** Non-steroidal anti-inflammatory, analgesic and antipyretic indicated for the treatment of pain and inflammation associated with canine osteoarthritis. / **SUPPLY AND DOSAGE:** Recommended dose is 2.2 mg./kg every 12 hours or 4.4 mg./kg every 24 hours. Length of the treatment is based on clinical opinion and according to the evolution of the clinical case. / **PRESENTATION:** Package of 60 tablets

- To control post-operative pain
- Selective inhibitor of COX2
- Rapid action
- Mininum gastric irritation

Prote-cis articular

TABLETS

Osteoarticular regenerator.
Chondrogenic, Anti-arthritis
for Canines and Felines



FORMULA: D. Glucosamine: 400 mg.; Chondroitin Sulfate: 100 mg.; Sodium Selenite: 5 mg.; Vitamin C: 30 mg.; Vitamin E: 50 mg.; Manganese Gluconate: 20 mg.; Excipients c.s. 1 tablet / **INSTRUCTIONS:** Osteoarthritis, primary and secondary arthritis, spondyloarthritis, discopathies, coxarthrosis due to dysplasia. To prevent disorders derived from hip dysplasia. It strengthens osteoarticular structures in growing animals. Anti-arthritis, joint analgesic. Antioxidant **SUPPLY AND DOSAGE:** 1 tablets per each 20 kg. liveweight during 60 days. Or better veterinary opinion. **PRESENTATION:** Package of 60 tablets.

- The most complete and best balanced FORMULA
- No contraindications

BIG ANIMALS

Aptomectin Fasciolida

ORAL SUSPENSION

FORMULA: Ivermectin: 0.2 gr.; Triclabendazole: 12 gr.; Excipients c.s. 100 ml. **INSTRUCTIONS:** For the treatment and control of endoparasitosis in ovines, to attack pulmonary nematodes, gastrointestinal, (including inhibited states of Ostertagia) For the treatment of hepatic fascioliasis (Gagayayae). / **SUPPLY AND DOSAGE:** Orally. 1 ml. per each 10 kg. liveweight. (200 mcg of Ivermectin per kg. liveweight and 12 mg of Triclabendazole per kg. liveweight). In case of severe infestations, dose can be repeated with an interval of not less than 120 days. / **PRESENTATION:** Drum of 2500 and 5000 ml.



AptoFen

INJECTABLE

FORMULA: Oxfendazole 12,5 g. / **INSTRUCTIONS:** For the treatment and control of gastrointestinal and pulmonary nematodes. Ovicide, larvicide and adulticide. Gastrointestinal nematodes: Ostertagia Ostertagi, Trichostrongylus Axei, Trichostrongylus C., Haemonchus Placei, Cooperia O., Oesophagostomum Radiatum, Oesophagostomum C. Pulmonary nematodes: Dictyocaulus Viviparus. / **SUPPLY AND DOSAGE:** Injectable subcutaneously, in front of or behind the pallet. 1ml. per each 50 kg liveweight (2,5 mg. per each kg. liveweight). Do not inject more than 10 ml per point of application. **PRESENTATION:** Bottle of 500 and 50 ml.



Afford



Afford Vademecum



HIGH QUALITY PRODUCTS
for animal health

Afford S.A.
Garibaldi 2619, CP (B1836ALH)
Parque Industrial Lomas de Zamora, Llavallol
Prov. de Buenos Aires, República Argentina - Telefax: (54 11) 4231.8155

www.afford.com.ar

f afford @afford_lab

